
Overview

This standard covers preparing for and performing intermediate life support, including the use of equipment to promote airway management and assist in ventilation. It addresses clearing and maintaining an open airway, establishing adequate ventilation and cardiac output in an individual as well as responsibility for monitoring and responding to changes in the individual's condition.

Users of this standard will need to ensure that practice reflects up to date information and policies.

SFHEC17

Provide intermediate life support

Performance criteria

You must be able to:

- P1 establish the need for intermediate life support based on best practice
- P2 check and confirm whether or not :
 - P2.1 there is a need to open or clear the airway of obstructions
 - P2.2 the individual is breathing unaided
 - P2.3 the individual's respiration is inadequate for life support
 - P2.4 the individual's circulation has stopped
 - P2.5 resuscitation is contraindicated
- P3 determine the appropriate equipment and techniques to use according to the individual's needs in order to:
 - P3.1 establish an open airway
 - P3.2 ventilate the individual
 - P3.3 establish and maintain cardiac output
- P4 clear the individual's airway of obstruction using the required technique, taking account of the individual's condition
- P5 position yourself and the individual appropriately to provide life support, taking due account of the individual's condition and circumstances
- P6 apply life support techniques and equipment consistent with:
 - P6.1 evidence based practice
 - P6.2 established, local protocols
 - P6.3 your own scope of practice
 - P6.4 legislation
- P7 seek additional support promptly where the individual's needs exceed your scope of practice
- P8 maintain adequate oxygenation and cardiac output for the individual until:
 - P8.1 adequate spontaneous breathing and cardiac output is established
 - P8.2 the individual is handed over to the care of others
 - P8.3 death is confirmed by an authorised practitioner
 - P8.4 you become exhausted
- P9 alternate compressions with ventilation of the lungs at the appropriate ratio
- P10 monitor and evaluate the effectiveness of your life support interventions, responding promptly and appropriately to achieve the best possible outcome for the individual
- P11 retain, for analysis, samples of any liquids blocking the oropharynx and mouth if appropriate to patient care
- P12 apply appropriate health, safety and infection control precautions throughout
- P13 reinstate equipment and materials to working status after use
- P14 complete all required documentation accurately and legibly

SFHEC17

Provide intermediate life support

Knowledge and understanding

You need to know and understand:

- K1 the legislation regarding confidentiality and information sharing, the provision of services, the rights of the individual, protection issues, anti-discriminatory practice, informed consent, relevant mental health legislation and care programme approach
- K2 the importance of gaining assent from individuals who lack capacity to consent
- K3 the ethics concerning consent and confidentiality, and the tensions which may exist between an individual's rights and the organisation's responsibility to individuals
- K4 how to interpret and apply legislation to the work being undertaken
- K5 the policy and procedures for summoning assistance for prolonged/extended resuscitation
- K6 the policy and procedures for recording information on the provision of ILS and the details which should be recorded
- K7 the anatomy and physiology of the respiratory system
- K8 the priorities in life support
- K9 the content of the Paediatric and Adult Intermediate Life Support Guidelines as issued by the UK Resuscitation Council
- K10 the time frame within which assessment of individual needs should be carried out and the life support response initiated
- K11 the differences in techniques needed for conducting advanced life support on different types of individual
- K12 why different techniques are needed depending on the individual's condition and circumstances
- K13 the factors to be taken into account in determining the technique that will lead to the best possible outcome for the individual, including contra-indications to the use of a given technique
- K14 the situations under which intermediate life support (ILS) should not be initiated, as set down in IAW and ILS guidelines (e.g. decapitation, submersion of an adult for longer than 30 minutes)
- K15 the measures to be taken to ensure health and safety, infection control and the prevention of contamination
- K16 the importance to outcome of the positioning of the individual and the practitioner applying intermediate life support (ILS), including the specific positioning needs of pregnant women in the third trimester
- K17 the anatomy and physiology relevant to different external chest compression techniques
- K18 the pulse sites and rate norms for different types of individual
- K19 the upper and lower readings for different types of individual on equipment used in monitoring clinical signs
- K20 the clinical signs and causes of airway obstruction

SFHEC17

Provide intermediate life support

- K21 the policies and procedures to be followed for the use of suction apparatus
- K22 what to do in the event of foreign body obstruction of an individual's airway
- K23 why the head tilt techniques should not be used where neck or spinal injury is suspected
- K24 the purpose of oropharyngeal and nasopharyngeal airways
- K25 the indications for the use of oropharyngeal and nasopharyngeal airways and the circumstances for their removal
- K26 the insertion procedures for oropharyngeal and nasopharyngeal airways
- K27 the procedures for suctioning in the presence of oropharyngeal and nasopharyngeal airways
- K28 the ventilation ratio and rate for different types of individual
- K29 the observations to be carried out to identify adequate oxygenation in different types of individual
- K30 the compression : ventilation ration in one and two person ILS
- K31 the procedure to establish the correct hand/finger placement for applying external chest compression
- K32 the procedures and equipment used for monitoring an individual's clinical signs during and post resuscitation
- K33 why adjustments to the techniques used in ILS may be needed, including the variations needed when the individual is hypothermic
- K34 different methods of chest thrusts and back slaps to use for children and young people, adults of a working age and older people
- K35 the handover procedures for an individual receiving ILS
- K36 the policies and procedures to be followed regarding samples of fluids suctioned from an individual

SFHEC17

Provide intermediate life support

Additional Information

External Links

This standard links with the following dimension within the NHS Knowledge and Skills Framework (October 2004):

Dimension: HWB7 Interventions and treatments

SFHEC17

Provide intermediate life support

Developed by	Skills for Health
Version number	1
Date approved	June 2010
Indicative review date	June 2012
Validity	Current
Status	Original
Originating organisation	Skills for Health
Original URN	EC17
Relevant occupations	Health, Public Services and Care; Nursing and Subjects and Vocations Allied; Healthcare and Related Personal Services
Suite	Emergency, Urgent and Scheduled Care
Key words	emergency care, scheduled care, urgent care